

Shelby County Preliminary Health Profile, Tennessee Adult Lifestyles Surveys, 1998

| Variable | Size | Count | Weighted Rate (%) | 95% CI | State Rate (%) | Rank Hi to Lo |
|---|------|-------|----------------------|--------------|-------------------|------------------|
| Q80.AGE 18-24 YEARS | 207 | 18 | 10.0 | (5.6, 14.3) | 12.0 | 60 |
| Q80.AGE 65+ YEARS | 207 | 38 | 12.5 | (8.4, 16.6) | 14.0 | 69 |
| Q116/Q98. BLACKS | 205 | 83 | 48.9* | (38.4, 59.5) | 14.1 | 1 |
| Q73. FEMALES | 207 | 134 | 53.5 | (45.4, 61.5) | 52.5 | 29 |
| Q113/Q95. HOUSEHOLD INCOME <= \$15,000 | 188 | 32 | 17.9 | (9.0, 26.9) | 17.0 | 59 |
| Q1 .IN FAIR/POOR HEALTH | 207 | 39 | 17.8 | (12.1, 23.4) | 17.3 | 63 |
| Q2 .OFTEN VERY DEPRESSED | 207 | 20 | 7.0 | (3.4, 10.6) | 8.9 | 80 |
| Q9 .HIGH BLOOD PRESSURE EVER | 207 | 66 | 28.6 | (21.6, 35.6) | 25.8 | 48 |
| Q9 .HIGH BLOOD PRESSURE IN PAST YR | 207 | 27 | 11.3 | (6.3, 16.4) | 10.8 | 57 |
| Q9 .HEART DISEASE EVER | 207 | 26 | 9.4 | (5.1, 13.7) | 10.1 | 74 |
| Q9 .HEART DISEASE IN PAST YR | 207 | 9 | 3.6 | (1.2, 5.9) | 3.4 | 52 |
| Q9 .DIABETES EVER | 207 | 21 | 10.5 | (4.8, 16.3) | 8.0 | 22 |
| Q9 .DIABETES IN PAST YR | 207 | 6 | 2.7 | (2.0, 3.5) | 2.5 | 48 |
| Q9 .STROKE EVER | 207 | 7 | 2.7 | (0.7, 4.6) | 1.9 | 35 |
| Q9 .STROKE IN PAST YR | 207 | 1 | 0.2 | (0.1, 0.5) | 0.3 | 49 |
| Q9 .SEIZURE EVER | 207 | 3 | 1.1* | (0.6, 1.6) | 2.2 | 74 |
| Q9 .SEIZURE IN PAST YR | 207 | 2 | 0.8 | (0.4, 1.2) | 0.6 | 40 |
| Q9 .LUNG DISEASE EVER | 207 | 12 | 3.5 | (1.4, 5.6) | 4.7 | 69 |
| Q9 .LUNG DISEASE IN PAST YR | 207 | 3 | 0.8 | (0.5, 1.4) | 1.5 | 63 |
| Q9 .PNEUMONIA EVER | 207 | 39 | 20.2 | (11.8, 28.7) | 19.0 | 42 |
| Q9 .PNEUMONIA IN PAST YR | 207 | 6 | 2.7 | (0.4, 4.9) | 2.6 | 44 |
| Q9 .TUBERCULOSIS EVER | 207 | 1 | 0.2 | (0.0, 0.5) | 0.5 | 65 |
| Q9 .CANCER EVER | 207 | 16 | 5.8 | (2.9, 8.8) | 6.2 | 55 |
| Q9 .CANCER IN PAST YR | 207 | 2 | 0.7 | (0.4, 1.2) | 1.4 | 65 |
| Q9 .CHRONIC STOMACH PROBLEMS EVER | 207 | 38 | 18.2 | (12.3, 24.0) | 17.5 | 57 |
| Q9 .CHRONIC STOMACH PROBLEMS IN PAST YR | 207 | 9 | 3.4 | (1.1, 5.8) | 5.9 | 82 |
| Q9 .GALLBLADDER PROBLEMS EVER | 207 | 18 | 4.8* | (2.5, 7.2) | 8.2 | 93 |
| Q9 .GALLBLADDER PROBLEMS IN PAST YR | 207 | 1 | 0.3* | (0.1, 0.6) | 1.1 | 80 |
| Q9 .LIVER PROBLEMS EVER | 207 | 3 | 0.8* | (0.4, 1.3) | 2.9 | 85 |
| Q9 .LIVER PROBLEMS IN PAST YR | 207 | 0 | 0.0 | (0.0, 0.2) | 0.3 | 40 |
| Q9 .INJURY EVER | 206 | 105 | 50.9 | (41.8, 60.0) | 57.3 | 80 |
| Q9 .INJURY IN PAST YR | 206 | 15 | 6.5 | (2.5, 10.6) | 9.3 | 77 |
| Q9 .HIV/AIDS EVER | 206 | 0 | 0.0 | (0.0, 0.2) | 0.1 | 15 |
| Q9 .JOINT PROBLEMS EVER | 207 | 57 | 27.1 | (19.0, 35.1) | 27.7 | 63 |
| Q9 .JOINT PROBLEMS IN PAST YR | 207 | 26 | 12.5 | (4.8, 20.2) | 12.2 | 48 |

*: There is a statistically significant difference (at 0.05 level) between the county and the state rate.

95% CI: 95% confidence interval.

Rank: Rankings go from highest to lowest rates. In all but 14 variables, higher rates are "worse."

Shelby County Preliminary Health Profile, Tennessee Adult Lifestyles Surveys, 1998

| Variable | Size | Count | Weighted Rate (%) | 95% CI | State Rate (%) | Rank Hi to Lo |
|--|------|-------|----------------------|--------------|-------------------|------------------|
| Q9 .KIDNEY DISEASE EVER | 207 | 8 | 3.2 | (1.0, 5.4) | 4.4 | 71 |
| Q9 .KIDNEY DISEASE IN PAST YR | 207 | 2 | 1.1 | (0.7, 1.7) | 1.1 | 42 |
| Q9 .MENTAL ILLNESS EVER | 201 | 12 | 4.2 | (1.2, 7.2) | 6.1 | 75 |
| Q9 .MENTAL ILLNESS IN PAST YR | 201 | 4 | 0.9* | (0.6, 1.5) | 2.5 | 84 |
| Q9 .LEARNING PROBLEMS EVER | 201 | 3 | 2.3 | (1.7, 3.1) | 2.6 | 49 |
| Q9 .LEARNING PROBLEMS IN PAST YR | 201 | 0 | 0.0 | (0.0, 0.2) | 0.4 | 42 |
| Q9 .OTHER MAJOR ILLNESS EVER | 207 | 31 | 10.0* | (6.1, 13.9) | 16.9 | 90 |
| Q9 .OTHER MAJOR ILLNESS IN PAST YR | 207 | 7 | 2.2* | (0.5, 3.9) | 4.5 | 82 |
| Q10.EVER INJURED IN TRAFFIC CRASH | 207 | 68 | 31.9 | (23.8, 39.9) | 32.1 | 40 |
| Q10.INJURED IN CRASH IN PAST YR | 207 | 6 | 2.2 | (0.4, 4.0) | 2.6 | 44 |
| Q10.INJURED IN CRASH IN PAST 30 DAYS | 206 | 1 | 0.3 | (0.1, 0.7) | 0.3 | 22 |
| Q11.INJURIES OTHER THAN FROM CRASHES | 207 | 121 | 57.5 | (49.9, 65.1) | 61.0 | 73 |
| Q11.OTHER INJURIES IN PAST YR | 207 | 14 | 7.2 | (2.8, 11.6) | 9.2 | 70 |
| Q11.OTHER INJURIES IN PAST 30 DAYS | 207 | 2 | 1.5 | (1.0, 2.2) | 1.7 | 48 |
| Q12.HAVE ANY KIND OF HEALTH CARE COVERAG | 207 | 194 | 92.7 | (88.2, 97.2) | 91.9 | 37 |
| Q12.PRIVATE INSURANCE PLAN | 207 | 132 | 65.3 | (56.2, 74.5) | 65.9 | 34 |
| Q12.TENNCARE | 207 | 22 | 12.4 | (4.7, 20.1) | 11.1 | 41 |
| Q12.MEDICARE | 207 | 34 | 15.4 | (7.7, 23.1) | 12.5 | 20 |
| Q12.OTHER TYPE OF INSURANCE | 206 | 6 | 3.2 | (0.5, 5.8) | 3.4 | 46 |
| Q12.SUPPLEMENTAL HEALTH INSURANCE | 193 | 53 | 26.0 | (16.9, 35.1) | 25.4 | 45 |
| Q13.COULD NOT GET MEDICAL CARE PAST YR | 207 | 23 | 14.4 | (6.4, 22.3) | 7.6 | 2 |
| Q14.HAS A USUAL SOURCE OF HEALTH CARE | 206 | 176 | 84.1 | (78.5, 89.8) | 85.2 | 65 |
| Q15.EVER SEEN A DOCTOR FOR PHYS HEALTH | 207 | 141 | 64.9 | (56.2, 73.6) | 70.6 | 75 |
| Q15.SAW DOCTOR FOR PHYS HEALTH PAST YR | 207 | 85 | 38.2 | (29.8, 46.7) | 37.1 | 41 |
| Q15.USED EMERG ROOM FOR CARE EVER | 207 | 136 | 64.2 | (56.3, 72.1) | 71.5 | 84 |
| Q15.USED EMERG ROOM FOR CARE IN PAST YR | 207 | 40 | 19.3 | (11.0, 27.6) | 19.6 | 50 |
| Q15.EVER ADMITTED TO HOSPITAL | 206 | 155 | 65.6* | (56.6, 74.7) | 76.2 | 95 |
| Q15.ADMITTED TO HOSPITAL IN PAST YR | 206 | 28 | 10.3 | (6.0, 14.6) | 10.0 | 55 |
| Q15.ROUTINE CHECKUP EVER | 206 | 187 | 89.0 | (83.9, 94.2) | 90.8 | 53 |
| Q15.ROUTINE CHECKUP IN PAST YR | 206 | 125 | 57.7 | (50.5, 64.9) | 54.6 | 21 |
| Q17.EVER RECEIVED MENTAL HEALTH TREATMNT | 206 | 31 | 11.0 | (6.1, 15.9) | 11.8 | 47 |
| Q17.MENTAL HEALTH RX IN PAST YR | 206 | 10 | 2.4 | (0.7, 4.1) | 3.7 | 63 |
| Q17.EVER ADMITTED FOR PSYCH PROBLEMS | 200 | 12 | 4.6 | (1.7, 7.6) | 3.1 | 18 |
| Q17.ADMITTED FOR PSYCH PROB IN PAST YR | 200 | 0 | 0.0 | (0.0, 0.2) | 0.3 | 29 |
| Q18.EVER ATTEMPTED SUICIDE | 207 | 5 | 1.2* | (0.1, 2.3) | 2.4 | 75 |

*: There is a statistically significant difference (at 0.05 level) between the county and the state rate.

95% CI: 95% confidence interval.

Rank: Rankings go from highest to lowest rates. In all but 14 variables, higher rates are "worse."

Shelby County Preliminary Health Profile, Tennessee Adult Lifestyles Surveys, 1998

| Variable | Size | Count | Weighted Rate (%) | 95% CI | State Rate (%) | Rank Hi to Lo |
|--|------|-------|----------------------|--------------|-------------------|------------------|
| Q18.SUICIDE ATTEMPT IN PAST YR | 207 | 0 | 0.0 | (0.0, 0.2) | 0.4 | 30 |
| Q20.EVER DRANK CAFFEINE | 207 | 204 | 97.6 | (96.8, 98.2) | 98.7 | 84 |
| Q20.DRANK CAFFEINE IN PAST YR | 207 | 199 | 95.7 | (92.1, 99.3) | 96.7 | 67 |
| Q20.DRANK CAFFEINE IN PAST 30 DAYS | 206 | 189 | 91.1 | (86.4, 95.8) | 94.1 | 90 |
| Q20.EVER HOOKED ON CAFFEINE | 204 | 54 | 22.2* | (15.7, 28.7) | 29.4 | 90 |
| Q20.NOW HOOKED ON CAFFEINE | 203 | 44 | 18.4 | (12.8, 24.1) | 23.5 | 85 |
| Q21.EVER SMOKED 100 CIGARETTES | 206 | 96 | 46.9 | (38.5, 55.3) | 51.1 | 86 |
| Q21.SMOKED IN PAST YR | 206 | 46 | 26.4 | (18.1, 34.7) | 30.1 | 77 |
| Q21.SMOKED IN PAST 30 DAYS | 206 | 44 | 25.3 | (16.9, 33.7) | 28.3 | 76 |
| Q22.EVER USED OTHER KINDS OF TOBACCO | 207 | 29 | 16.7 | (8.7, 24.7) | 23.9 | 94 |
| Q22.USED OTHER TOBACCO IN PAST YR | 207 | 9 | 4.6* | (1.4, 7.7) | 11.5 | 95 |
| Q22.USED OTHER TOBACCO IN PAST 30 DAYS | 207 | 7 | 3.6* | (0.7, 6.4) | 8.4 | 94 |
| Q23.EVER HOOKED ON TOBACCO | 203 | 57 | 31.4 | (22.8, 40.0) | 37.2 | 88 |
| Q23.NOW HOOKED ON TOBACCO | 202 | 34 | 21.9 | (13.6, 30.2) | 25.3 | 83 |
| Q24.EVER DRANK ALCOHOL | 207 | 164 | 79.8 | (73.6, 85.9) | 79.5 | 33 |
| Q24.DRANK ALCOHOL PAST YR | 207 | 114 | 57.2 | (49.0, 65.3) | 52.5 | 12 |
| Q24.BINGED PAST YR (5+DRINKS) | 207 | 22 | 11.5 | (6.5, 16.6) | 15.8 | 79 |
| Q24.DRANK ALCOHOL PAST 30 DAYS | 207 | 83 | 42.8 | (35.2, 50.4) | 35.9 | 6 |
| Q24.DRANK ALCOHOL PAST 2 WEEKS | 207 | 64 | 30.5 | (24.1, 37.0) | 28.1 | 19 |
| Q24.EVER HOOKED ON ALCOHOL | 207 | 16 | 8.9 | (4.2, 13.6) | 6.7 | 19 |
| Q24.NOW HOOKED ON ALCOHOL | 205 | 1 | 0.5 | (0.2, 0.9) | 1.0 | 65 |
| Q25.EVER USED SEDATIVES | 205 | 33 | 12.1 | (7.4, 16.7) | 15.8 | 79 |
| Q25.USED SEDATIVES IN PAST YR | 205 | 11 | 3.5* | (0.9, 6.1) | 6.4 | 86 |
| Q25.USED SEDATIVES IN PAST 30 DAYS | 205 | 5 | 1.3* | (0.1, 2.6) | 3.5 | 90 |
| Q25.EVER USED TRANQUILIZERS | 207 | 48 | 19.6 | (11.8, 27.5) | 26.7 | 84 |
| Q25.USED TRANQUILIZERS IN PAST YR | 207 | 19 | 6.1* | (2.4, 9.8) | 10.7 | 89 |
| Q25.USED TRANQUILIZERS IN PAST 30 DAYS | 207 | 19 | 6.1 | (2.4, 9.8) | 7.4 | 62 |
| Q25.EVER USED STIMULANTS | 207 | 20 | 8.2 | (4.5, 12.0) | 9.5 | 64 |
| Q25.USED STIMULANTS IN PAST YR | 207 | 1 | 0.2* | (0.1, 0.5) | 1.4 | 80 |
| Q25.USED STIMULANTS IN PAST 30 DAYS | 207 | 0 | 0.0* | (0.0, 0.2) | 0.6 | 53 |
| Q25.EVER USED PAINKILLERS | 205 | 116 | 57.3 | (48.7, 65.8) | 60.2 | 64 |
| Q25.USED PAINKILLERS IN PAST YR | 205 | 45 | 25.4 | (16.4, 34.3) | 22.2 | 25 |
| Q25.USED PAINKILLERS IN PAST 30 DAYS | 205 | 20 | 8.3 | (4.1, 12.6) | 8.4 | 53 |
| Q26.EVER ABUSED PRESCRIPTION DRUGS | 207 | 5 | 2.3* | (0.1, 4.5) | 6.5 | 85 |
| Q26.ABUSED PRESCRIP DRUGS IN PAST YR | 207 | 0 | 0.0* | (0.0, 0.2) | 1.2 | 59 |

*: There is a statistically significant difference (at 0.05 level) between the county and the state rate.

95% CI: 95% confidence interval.

Rank: Rankings go from highest to lowest rates. In all but 14 variables, higher rates are "worse."

Shelby County Preliminary Health Profile, Tennessee Adult Lifestyles Surveys, 1998

| Variable | Size | Count | Weighted Rate (%) | 95% CI | State Rate (%) | Rank Hi to Lo |
|--|------|-------|----------------------|--------------|-------------------|------------------|
| Q26.ABUSED PRESCRIP DRUGS IN PAST 30 DAY | 207 | 0 | 0.0 | (0.0, 0.2) | 0.3 | 32 |
| Q27.EVER HOOKED ON PRESCRIPTION DRUGS | 207 | 0 | 0.0* | (0.0, 0.2) | 2.0 | 83 |
| Q27.NOW HOOKED ON PRESCRIPTION DRUGS | 207 | 0 | 0.0* | (0.0, 0.2) | 0.4 | 41 |
| Q28.EVER USED STEROIDS | 205 | 31 | 10.7* | (6.8, 14.6) | 15.7 | 79 |
| Q28.USED STEROIDS IN PAST YR | 205 | 15 | 4.3 | (2.2, 6.5) | 5.6 | 61 |
| Q28.USED STEROIDS IN PAST 30 DAYS | 205 | 6 | 1.5 | (0.3, 2.7) | 2.2 | 61 |
| Q28.EVER ABUSED STEROIDS | 205 | 0 | 0.0* | (0.0, 0.2) | 0.4 | 28 |
| Q28.EVER HOOKED ON STEROIDS | 205 | 0 | 0.0 | (0.0, 0.2) | 0.2 | 14 |
| Q28.NOW HOOKED ON STEROIDS | 205 | 0 | 0.0 | (0.0, 0.2) | 0.1 | 8 |
| Q29.EVER USED MARIJUANA | 207 | 55 | 30.7 | (22.7, 38.6) | 28.8 | 19 |
| Q29.USED MARIJUANA IN PAST YR | 207 | 7 | 6.9 | (5.8, 8.1) | 5.1 | 17 |
| Q29.USED MARIJUANA IN PAST 30 DAYS | 207 | 4 | 2.4 | (1.8, 3.2) | 2.6 | 39 |
| Q29.EVER HOOKED ON MARIJUANA | 207 | 3 | 2.4 | (1.7, 3.2) | 2.2 | 31 |
| Q29.NOW HOOKED ON MARIJUANA | 207 | 1 | 0.3 | (0.1, 0.8) | 0.5 | 33 |
| Q30.EVER USED COCAINE | 207 | 10 | 4.5 | (1.8, 7.2) | 7.0 | 68 |
| Q30.USED COCAINE IN PAST YR | 207 | 0 | 0.0* | (0.0, 0.2) | 0.8 | 46 |
| Q30.USED COCAINE IN PAST 30 DAYS | 207 | 0 | 0.0 | (0.0, 0.2) | 0.2 | 22 |
| Q30.EVER HOOKED ON COCAINE | 207 | 0 | 0.0* | (0.0, 0.2) | 0.8 | 54 |
| Q30.NOW HOOKED ON COCAINE | 207 | 0 | 0.0 | (0.0, 0.2) | 0.1 | 11 |
| Q31.EVER USED HEROIN | 207 | 2 | 1.0 | (0.6, 1.5) | 0.9 | 28 |
| Q31.USED HEROIN IN PAST YR | 207 | 0 | 0.0 | (0.0, 0.2) | 0.1 | 5 |
| Q31.EVER HOOKED ON HEROIN | 207 | 0 | 0.0 | (0.0, 0.2) | 0.1 | 12 |
| Q32.EVER USED HALLUCINOGENS | 207 | 6 | 1.9* | (0.1, 3.6) | 5.7 | 86 |
| Q32.USED HALLUCINOGENS IN PAST YR | 207 | 1 | 0.3 | (0.1, 0.8) | 0.8 | 29 |
| Q32.USED HALLUCINOGENS IN PAST 30 DAYS | 207 | 0 | 0.0 | (0.0, 0.2) | 0.2 | 11 |
| Q32.EVER HOOKED ON HALLUCINOGENS | 206 | 0 | 0.0 | (0.0, 0.2) | 0.1 | 11 |
| Q33.EVER USED INHALANTS | 207 | 4 | 1.7 | (1.2, 2.4) | 2.1 | 41 |
| Q33.USED INHALANTS IN PAST YR | 207 | 0 | 0.0 | (0.0, 0.2) | 0.2 | 14 |
| Q33.EVER HOOKED ON INHALANTS | 207 | 0 | 0.0 | (0.0, 0.2) | 0.2 | 14 |
| Q33.NOW HOOKED ON INHALANTS | 207 | 0 | 0.0 | (0.0, 0.2) | 0.1 | 4 |
| Q34.EVER INJECTED DRUG FOR NON-MED REASO | 207 | 1 | 0.3 | (0.1, 0.7) | 0.7 | 48 |
| Q35.EVER DRIVEN AFTER DRINKING | 207 | 111 | 56.5 | (48.7, 64.4) | 49.4 | 7 |
| Q35.DRIVEN AFTER DRINKING IN PAST YR | 207 | 34 | 18.1 | (11.9, 24.3) | 12.9 | 5 |
| Q35.EVER DRIVEN AFTER 5 OR MORE DRINKS | 207 | 48 | 25.4 | (18.1, 32.7) | 24.6 | 49 |
| Q35.DRIVEN AFTER 5 DRINKS IN PAST YR | 207 | 10 | 5.7 | (2.2, 9.2) | 3.4 | 12 |

*: There is a statistically significant difference (at 0.05 level) between the county and the state rate.

95% CI: 95% confidence interval.

Rank: Rankings go from highest to lowest rates. In all but 14 variables, higher rates are "worse."

Shelby County Preliminary Health Profile, Tennessee Adult Lifestyles Surveys, 1998

| Variable | Size | Count | Weighted Rate (%) | 95% CI | State Rate (%) | Rank Hi to Lo |
|--|------|-------|----------------------|--------------|-------------------|------------------|
| Q35.EVER DRIVEN AFTER USING PRESCRIP DRU | 207 | 75 | 32.4 | (24.7, 40.1) | 33.2 | 53 |
| Q35.DRIVEN AFTER PRESCRIPTION IN PAST YR | 207 | 48 | 21.7 | (15.0, 28.4) | 18.1 | 21 |
| Q35.EVER DRIVEN AFTER USING OTHER DRUGS | 207 | 22 | 8.8 | (5.1, 12.4) | 10.7 | 55 |
| Q35.DRIVEN AFTER OTHER DRUGS IN PAST YR | 207 | 6 | 2.2 | (1.6, 3.0) | 3.1 | 52 |
| Q67.EVER RECEIVED RX FOR ALCOHOL PROBLEM | 207 | 3 | 1.3* | (0.9, 1.9) | 2.5 | 70 |
| Q67.RECEIVED RX FOR ALCOHOL PROB IN PAST | 207 | 1 | 0.3 | (0.1, 0.7) | 0.4 | 41 |
| Q68.EVER RECEIVED RX FOR OTHER DRUG PROB | 207 | 0 | 0.0* | (0.0, 0.2) | 1.1 | 59 |
| Q68.RX FOR OTHER DRUG PROB IN PAST YR | 207 | 0 | 0.0 | (0.0, 0.2) | 0.2 | 17 |

Source: Community Health Research Group at the University of Tennessee, Knoxville, and Tennessee Department of Health

*: There is a statistically significant difference (at 0.05 level) between the county and the state rate.

95% CI: 95% confidence interval.

Rank: Rankings go from highest to lowest rates. In all but 14 variables, higher rates are "worse."